

Shark cartilage extracts as antiangiogenic agents: smart drinks or bitter pills?

Denis Gingras, Alain Renaud, Nathalie Mousseau and Richard Béliveau
*Laboratoire de médecine moléculaire Hôpital Ste-Justine-UQAM, Centre de cancérologie
Charles-Bruneau, Centre de Recherche de l'Hôpital Ste-Justine, Chemin Côte-Ste-Catherine, Montréal,
Québec, Canada*

Key words: shark cartilage, Æ-941, angiogenesis, matrix metalloproteinase, cancer

Abstract

The use of crude cartilage for the treatment of human cancers remains a subject of controversy. In this brief commentary, we reviewed the current knowledge on the anticancer properties of cartilage. We then presented the properties of Æ-941, a novel standardized water-soluble extract derived from shark cartilage that represents less than 5% of the crude cartilage. It is a multifunctional antiangiogenic product that contains several biologically active molecules. Æ-941 is one of the few antiangiogenic drugs that is under phase III clinical investigation. It is currently evaluated in Europe and North America for the treatment of refractory renal cell carcinoma and in North America for metastatic non small cell lung cancer.

Abbreviations: MMP – matrix metalloproteinase; TIMP – tissue inhibitor of metalloproteinase; VEGF – vascular endothelial growth factor.

Controversy and scientific reality

There is still considerable controversy regarding the clinical usefulness of cartilage for the treatment of cancer. Most of this controversy arises from the false claims that sharks (elasmobranch family) which skeleton is made by cartilage (about 6% of the whole body weight) do not develop cancers and that solid forms of shark cartilage may cure cancer in animals and humans. As a result, a large variety of rather expensive powders from total preparations of cartilage have invaded the natural product market. Over the last 25 years, a number of studies have claimed that powdered cartilage extracts were efficient for the treatment of cancer patients but most of these studies were received with scepticism by the scientific community due to their lack of adequate protocols. In the rare cases where these studies were conducted with a significant number of patients, no consistent improvement of patient conditions could be observed [1]. A recent Phases I/II clinical trial on cancer patients failed to support a beneficial effect of shark cartilage powder [2]. Moreover, the precise composition

of these crude preparations are unknown and the biological activities supporting the use of such products in the cure of solid tumors and other related diseases have never been established.

On the other hand, cartilage from all species is an avascular tissue known to develop malignant tumors only in rare occasions [3–6]. This resistance of cartilage to tumor formation seems to be correlated to the capacity of the tissue to inhibit the formation of new blood vessels which are necessary for the growth and the metastatic power of tumor cells [7–9]. In a series of experiments, Brem and Folkman [10] demonstrated that fractions or liquid extracts of cartilage were able to inhibit the proliferation of capillaries induced by tumor cells. Subsequently, it was shown that a cartilage fraction extracted by guanidium and purified by affinity chromatography was also able to inhibit angiogenesis induced by tumor cells [11,12]. This property of cartilage is conserved throughout evolution as antiangiogenic activities were observed in several species such as shark [13–16], bovine [10], and human [17]. Although the mechanisms involved in these antitumor

properties of cartilage extracts are not understood in detail, the purification of some of cartilage antitumor factors have suggested the potential involvement of anti-collagenase activities in their effects [16,18–22]. The antiangiogenic properties of bovine cartilage extracts were mainly correlated with the presence of at least two distinct type I collagenase inhibitors, named CDI and BCDI [18,23], that share extensive homologies with TIMP-1 and TIMP-2, respectively. This has led to the suggestion that these MMP inhibitors may account for many of the antiangiogenic properties of cartilage and may thus represent attractive therapeutic anticancer agents [19]. In the last decade, other cartilage-associated proteins with antiangiogenic properties such as chondromodulin [24], troponin I [25] and thrombospondin I [26] have been identified, further emphasizing the potential use of cartilage as an abundant source of inhibitors of neovascularization and therefore of potential anticancer drugs. However, the therapeutic potential of cartilage extracts relies on the adequate extraction of these molecules as well as on the preservation of their biological activities following their extraction and storage. In this respect, it is likely that commercial shark cartilage powders that contain a high proportion of inorganic material may have a low content in these inhibitors that would explain their low efficacy.

Æ-941, a novel water-soluble shark cartilage extract with antiangiogenic properties

Æ-941 is a standardized shark cartilage extract produced and tested according to CGMP criteria. Æ-941 differs from the previously reported solid extracts since it is a water-soluble extract from which more than 95% of inactive molecules have been removed, resulting in high concentrations of biologically-relevant molecules [27]. Following extraction, soluble molecules are separated from the solid part by centrifugation, and ultrafiltration is performed with a cut-off of 500 kDa to obtain the liquid extract, the Æ-941, and the extract is kept frozen until use to maximally preserve its biological properties. This mode of extraction thus differs considerably from that of other commercial preparations and may explain the preservation of the antiangiogenic properties of Æ-941. In particular, it was demonstrated that the compound has a marked inhibitory effect on embryonic vascularization (Figure 1), endothelial cell proliferation [28,29] and tubulogenesis [30]. The antiangiogenic activity of Æ-941 was correlated with the presence of an antimetalloproteinase inhibitor that inhibits several members of the MMP family, with a preferential inhibition of MMP-2 (Figure 2A) as well as with that of a serine elastase inhibitory activity (Figure 2B). In addition,




Figure 1. Effect of Æ-941 on embryonic neovascularization. (A) *Ex ovo* chick embryo culture were treated with methylcellulose disk containing either a control solution or 200 µg Æ-941. Discs were deposited on the external border of the vascular perimeter near the tip of blood vessels, where the angiogenic process is active. In order to minimize inter-individual variations, discs containing either the drug sample or the vehicle were placed in a symmetric fashion with respect to the cephalo-caudal axis of the embryo, since blood vessels develop symmetrically to that axis. *Ex ovo* incubation of the embryos was pursued for 24 h. Ten chick embryos per conditions were examined. (B) Increasing concentration of Æ-941 were added to the chick embryo and angiogenesis was assessed in a blind semi-quantitative fashion. The blood vessel formation was considered as altered when its growing path was either deviated, diminished or when there was no growth observed beyond the disc as compared to the negative control.




Figure 2. Effect of Æ-941 on the gelatinolytic and elastinolytic activities of matrix metalloproteinase-2 and serine elastases. Recombinant MMP-2 (A) or porcine pancreatic elastase (PPE) (B) were incubated in the absence (dark circles) or in the presence (open circles) of 100 µg/ml of Æ-941 in a reaction buffer containing gelatin or elastin labeled with BODIPY FL. The proteolytic activities of both enzymes were determined by following the increase in fluorescence at 538 nm. Under these conditions, Æ-941 inhibited MMP-2 and PPE activities by 60% and 98%, respectively.

Æ-941 inhibits VEGF binding to endothelial cells [31], the VEGF-dependent tyrosine phosphorylation of the VEGF receptor as well as the VEGF-dependent increase in vascular permeability [30].

In vivo, Æ-941 was also found to have significant *in vivo* antitumor and antimetastatic activities [32]. In a therapeutic point of view, recent results from a phases I–II clinical trial in patients suffering from a refractory non-small cell lung cancer revealed that Æ-941 is orally bioavailable and safe, since no serious side effects related to the administration of Æ-941 have been observed. Furthermore, a retrospective analysis

indicates that Æ-941 significantly increase the median survival of these patients [33]. Æ-941 is one of the few antiangiogenic drugs that has reached phase III clinical trial evaluation. It is currently tested in phase III clinical trials in Europe and North America for the treatment of refractory renal cell carcinoma and in North America for metastatic lung cancers.

Conclusions

Scientific data obtained from various studies where Æ-941 has been tested strongly suggest that controlled water extraction and frozen storage of cartilage represent crucial steps in order to retain its multiple biological activities. By contrast to powdered shark cartilage preparations that show no significant effects on cancer patient's conditions, Æ-941 has been used in well-controlled clinical settings and emerged as a compound that can significantly improve survival. Given the recent studies supporting the presence within Æ-941 of specific inhibitor(s) of VEGF-mediated signal transduction events together with proteinases inhibitors, this compound appears as a multitarget drug contrasting from other natural or chemical antiangiogenic agents by its efficiency against several crucial steps of the angiogenic cascade closely linked to tumor progression. Further characterization and purification of the molecular components that are responsible of the biological activity of Æ-941, including the inhibition of specific matrix-degrading enzymes, serine elastases, and the control of endothelial cell proliferation will be of significant importance in our understanding of the molecular and cellular mechanisms involved in the antiangiogenic and antimetastatic properties of Æ-941.

References

1. Prudden JF: The treatment of human cancer with agents prepared from bovine cartilage. *J Biol Response Mod* 4: 551–584, 1985
2. Miller DR, Anderson GT, Stark JJ, Granick JL, Richardson D: Phase I/II trial of the safety and efficacy of shark cartilage in the treatment of advanced cancer. *J Clin Oncol* 16: 3649–3655, 1998
3. Eisenstein R, Sorgente N, Soble LW, Miller AW, Kuettner KE: The resistance of certain tissues to invasion: penetrability of explanted tissues by vascularized mesenchyme. *Am J Physiol* 73: 765–774, 1973
4. Eisenstein R, Kuettner KE, Neopolitan C, Soble LW, Sorgente N: The resistance of certain tissue to invasion. III. Cartilage extracts inhibit the growth of fibroblasts and endothelial cells in culture. *Am J Pathol* 81: 337–347, 1975

5. Harshbarger JC, Ostrander GK: Cancer in sharks, skates, rays and other lower fishes. In: American Association for Cancer Research, San Francisco, 2000, 762 pp.
6. Sorgente N, Kuettner KE, Soble LW, Eisenstein R: The resistance of certain tissues to invasion. II. Evidence of extractable factors in cartilage which inhibit invasion by vascularized mesenchyme. *Lab Invest* 32: 217–222, 1975
7. Folkman J: Tumor angiogenesis: therapeutic implications. *New Engl J Med* 285: 1182–1186, 1971
8. Folkman J: Anti-angiogenesis: new concept for therapy of solid tumors. *Ann Surgery* 175: 409–416, 1972
9. Folkman J: Tumor angiogenesis. *Adv Cancer Res* 43: 175–203, 1985
10. Brem H, Folkman J: Inhibition of tumor angiogenesis mediated by cartilage. *J Exp Med* 141: 427–439, 1975
11. Langer R, Brem H, Faltermann K, Klein M, Folkman J: Isolation of a cartilage factor that inhibits tumor neovascularization. *Science* 193: 70–72, 1976
12. Langer R, Conn H, Vacanti J, Haudenschild C, Folkman J: Control of tumor growth in animals by infusion of an angiogenesis inhibitor. *Proc Natl Acad Sci USA* 77: 4331–4335, 1980
13. Lee A, Langer R: Shark cartilage contains inhibitors of tumor angiogenesis. *Science* 221: 1185–1187, 1983
14. McGuire TR, Kazakoff PW, Hoie EB, Fienhold MA: Antiproliferative activity of shark cartilage with and without tumor necrosis factor- α in human umbilical vein endothelium. *Pharmacotherapy* 16: 237–244, 1996
15. Oikawa T, Ashino-Fuse H, Shimamura M, Koide U, Iwaguchi T: A novel angiogenic inhibitor derived from Japanese shark cartilage (I). Extraction and estimation of inhibitory activities toward tumor and embryonic angiogenesis. *Cancer Lett* 51: 181–186, 1990
16. Sheu JR, Fu CC, Tsai ML, Chung WJ: Effect of U-995, a potent shark cartilage-derived angiogenesis inhibitor, on anti-angiogenesis and antitumor activities. *Anticancer Research* 18: 4435–4442, 1998
17. Kaminski M, Kaminska G, Majewski S: Inhibition of new blood vessel formation in mice by systemic administration of human rib cartilage extract. *Experientia* 34: 490–491, 1978
18. Moses MA, Sudhalter J, Langer R: Identification of an inhibitor of neovascularization from cartilage. *Science* 248: 1408–1410, 1990
19. Moses MA, Langer RA: A metalloproteinase inhibitor as an inhibitor of neovascularization. *J Cell Biochem* 47: 230–235, 1991
20. Moses MA, Sudhalter J, Langer R: Isolation and characterization of an inhibitor of neovascularization from scapular chondrocytes. *J Cell Biol* 119: 475–482, 1992
21. Ohba Y, Goto Y, Kimura Y, Suzuki F, Hisa T, Takahashi K, Takigawa M: Purification of an angiogenesis inhibitor from culture medium conditioned by a human chondrosarcoma-derived chondrocytic cell line, HCS-2/8. *Biochim Biophys Acta* 1245: 1–8, 1995
22. Kuettner KE, Soble L, Croxen RL, Marzyska B, Hiti J, Harper E: Tumor cell collagenase and its inhibition by a cartilage derived protease inhibitor. *Science* 196: 653–654, 1977
23. Murray JB, Allison K, Sudhalter J, Langer R: Purification and partial amino acid sequence of bovine cartilage-derived collagenase inhibitor. *J Biol Chem* 261: 4154–4159, 1986
24. Hiraki Y, Tanaka H, Inoue H, Kondo J, Kamizono A, Suzuki F: Molecular cloning of a new class of cartilage-specific matrix, chondromodulin-I, which stimulates growth of cultures chondrocytes. *Biochem Biophys Res Commun* 175: 971–975, 1991
25. Moses MA, Wiederschain D, Wu I, Fernandez CA, Ghazizadeh V, Lane WS, Flynn E, Sytkowski A, Tao T, Langer R: Troponin I is present in human cartilage and inhibits angiogenesis. *Proc Natl Acad Sci USA* 96: 2645–2650, 1999
26. DiCesare PE, Morgelin M, Mann K, Paulsson M: Cartilage oligomeric matrix protein and thrombospondin 1. Purification from articular cartilage, electron microscopic structure, and chondrocyte binding. *Eur J Biochem* 223: 927–937, 1994
27. Dupont E, Brazeau P, Juneau C: Extracts of shark cartilage having an antiangiogenic activity and an effect on tumor progression: process of making thereof. United States Patent, #5,618,925, 1995
28. Dupont E, Savard PE, Thibodeau A, Vincent M, Chamberland G, Doillon C: Angiostatic activity of Neovastat (\mathcal{A} -941) assessed by a semi-quantitative embryonic vascularization test. In: IBC's Conference. Therapeutic Implications and Mechanisms of Angiogenesis, Novel Therapeutic Development, Boston, US, 1997 (Abstract)
29. Dupont E, Alaoui-Jamali M, Wang T, Thibodeau A, Alpert L, Savard P, Falardeau P: *In vitro* and *in vivo* anti-tumor activity of Neovastat (\mathcal{A} -941), an angiostatic agent derived from shark cartilage. In: IBC's Conference. Therapeutic Implications and Mechanisms of Angiogenesis, Novel Therapeutic Development, Boston, US, 1997 (Abstract)
30. Beliveau R, Sirois P, Tranqui L, Baffert F, Dimitriadou V, Pepin MC, Sirois MG, Poyet P, Falardeau P, Dupont E: Antiangiogenic agent \mathcal{A} -941 induces *in vivo* and *in vitro* inhibition of VEGF. In: American Association for Cancer Research, San Francisco, 2000 (Abstract)
31. Sirois MG: \mathcal{A} -941, a new and specific VEGF antagonist? In: American Association for Cancer Research, Philadelphia, US, April, 10–14, 1999 (Abstract)
32. Rivière M, Alaoui-Jamali M, Falardeau P, Blasecki J, Dupont E: \mathcal{A} -941 (Neovastat): an inhibitor of angiogenesis with anti-cancer activity. In: American Association for Cancer Research, New Orleans, US, 1998 (Abstract)
33. Latreille J, Laberge F, Rivière M, Falardeau P, Champagne P, Croteau D, Dupont E, Levinton C: Phase I/II clinical trials of escalating dose of \mathcal{A} -941, an inhibitor of angiogenesis in patients with refractory lung cancer. In: Proceedings of the 90th Annual Meeting of the American Association for Cancer Research, Philadelphia, US, 1999 (Abstract)

Address for offprints: Richard Béliveau, Laboratoire de médecine moléculaire Ste-Justine-UQAM, Centre de cancérologie Charles-Bruneau, 3175, Chemin Côte-Ste-Catherine, Montréal, Québec Canada H3T 1C5; *Fax:* (514)345-4731; *e-mail:* molmed@justine.umontreal.ca